

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11) Publication number:

0 408 965 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification: **04.10.95** (51) Int. Cl.⁶: **C11D 1/825**, C11D 1/83,

(21) Application number: **90112587.2**

C11D 1/94, A61K 7/08,
A61K 7/50

(22) Date of filing: **02.07.90**

(54) **Neutral liquid detergent composition.**

(30) Priority: **18.07.89 JP 184980/89**

(43) Date of publication of application:
23.01.91 Bulletin 91/04

(45) Publication of the grant of the patent:
04.10.95 Bulletin 95/40

(84) Designated Contracting States:
DE ES FR GB

(56) References cited:
EP-A- 0 075 995
EP-A- 0 105 556
FR-A- 2 358 877
US-A- 4 668 422

(73) Proprietor: **KAO CORPORATION**
1-14-10, Nihonbashikayaba-cho
Chuo-ku
Tokyo (JP)

(72) Inventor: **Deguchi, Katsuhiko**
2-4-12-507, Izumigaoka
Utsunomiya-shi,
Tochigi (JP)
Inventor: **Saito, Kozo**
2-4-12-203, Izumigaoka
Utsunomiya-shi,
Tochigi (JP)
Inventor: **Saijo, Hiroyuki**
152 -39, Miyukigahara-machi
Utsunomiya-shi,
Tochigi (JP)

(74) Representative: **Wächtershäuser, Günter, Prof.**
Dr.
Patentanwalt,
Tal 29
D-80331 München (DE)

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

EP 0 408 965 B1

Description**BACKGROUND OF THE INVENTION**5 **Field of the Invention:**

The present invention relates to a detergent composition, and, more particularly, to a neutral liquid detergent composition exhibiting remarkably reduced irritation and damage to the hair or the skin, possessing high foaming ability, and imparting a favorable feeling upon use. The detergent composition has also good rinsability of foam and excellent drainability from tableware and the like after washing.

Description of the Background Art:

Because of increased awareness for safety towards the human body in recent years, a number of trials have been undertaken in order to moderate the action of detergent compositions which frequently contact the skin. Such detergent compositions include laundry detergents, dishwashing detergents, household detergents, hair shampoos, and body shampoos. One of the approaches is adjusting the pH of detergent compositions to a weakly acidic range, e.g. pH 5-6, close to the intrinsic pH of human skins so as to be mild to the skin. Another approach is to incorporate low irritant components as major detergent base components. For example, amino acid surfactants, alkyl phosphate surfactants, or the like are being used as a low irritant detergent (Japanese Patent Publication Nos. 40125/1975, 426023/1976, 9033/1980, and 27319/1983).

Even though these surfactants are low irritant, they do not necessarily exhibit sufficient detergency and foaming ability is also their drawback. On the other hand, sodium alkylbenzenesulfonates have conventionally been popular detergent base components for dishwashing detergents because of their strong detergency. The compounds, however, give a strong defatting action to the skin and thus can be a cause of roughened hands.

Because of these reasons, sodium alkylethoxysulfates which are less irritant to the skin have gained a wider popularity as the base components for dishwashing detergents in recent years. Their detergent performances are even promoted by incorporating co-surfactants such as tertiary alkyl amine oxides, higher fatty acid diethanolamides, and the like. Their mildness to the skin is also improved by such a combination as compared with conventional detergent compositions.

The mildness to the skin of the detergents has improved to some degree as described above, nevertheless the achievement at present is still unsatisfactory.

Alkyl glycosides, being saccharide-derived surface active agents, are less irritant compounds. Although they are nonionic surfactants, it is well known that they not only produce stable foam themselves but also can act as a foam stabilizer for anionic surfactants. Thus, alkyl glycosides are gaining much attention in recent years. EP-A-0 105 556 and EP-A-0 075 995 disclose detergent compositions containing alkylpolysaccharides and nonionic surfactants.


Japanese Patent Laid-open No. 104625/1983, for example, discloses a foaming surfactant composition comprising an alkyl glycoside and an anionic surface active agent, and Japanese Patent Laid-open No. 74999/1987 discloses a liquid dishwashing detergent composition comprising an alkyl glycoside, an anionic surface active agent, and a fatty acid alkanolamide. They claim that the compositions are low irritant and have an excellent detergency. Even though these detergent compositions exhibit performances better than conventional detergent compositions containing polyoxyethylenealkylethers as a base component, such performances are still to be improved. They have problems such as an inferior feeling upon use, poor rinsability of the foam, and insufficient drainability.

In view of this situation, the present inventors have conducted extensive studies to make the maximum use of the characteristics of alkyl glycosides, and found that a detergent composition which exhibits an excellent feeling upon use, good rinsability of the foam, and superior drainability by incorporating alkyl glycosides together with a specific amount of certain type of nonionic surfactant having a specific HLB other than alkyl glycosides and a specific amount of certain type of water-soluble salt. Such findings have led to the completion of this invention.

55 **SUMMARY OF THE INVENTION**

Accordingly, an object of the present invention is to provide a neutral liquid detergent composition comprising:

(a) 2-60% by weight of an alkyl glycoside represented by the following formula (I):



5 wherein R_1 is a linear or branched alkyl, alkenyl, or alkylphenyl group having 8-18 carbon atoms, R_2 is an alkylene group having 2-4 carbon atoms, G is a reducing saccharide residue having 5-6 carbon atoms, x is a number of which the mean value is 0-5, and y is a number of which the mean value is 1.0-1.42;

(b) 0.1-10% by weight of a nonionic surfactant having an HLB of less than 5;

10 (c) 0.1-10% by weight of a nonionic surfactant having an HLB of not less than 5; and

(d) 0.001-8% by weight of one or more water-soluble organic or inorganic salts selected from the group consisting of sulfates, chlorides, borates, phosphates, p-toluenesulfonates, m-xylenesulfonates, benzoates, malates, succinates, tartarates, citrates, lactates, and edates of sodium and potassium; the ratio by weight of components [(b)+(c)]/(a) being in a range of 1/10-1/1 and the ratio by weight of components (b)/(c) being in a range of 1/10-1/1.

15 Other objects, features and advantages of the invention will hereinafter become more readily apparent from the following description.

DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS

20 Alkyl glycoside, component (a) of the present invention, is used as a major surfactant in the composition of the present invention. The mean value of x in formula (I), being defined as 0-5, affects the water-solubility and crystalline property of the alkyl glycosides. Higher the value x, are the alkyl glycosides more soluble in water and less crystalline. A desirable value of x is in the range of 0-2.

25 When the value y in formula (I) is larger than 1, i.e., when alkyl glycosides contain those having di- or higher saccharides as the hydrophilic groups, the saccharide chain bonds may be 1-2, 1-3, 1-4, 1-6, α -pyranoside, β -pyranoside, or franoside, or a mixture thereof. Among the preferable value of y which is 1.0-1.42, a most preferable range is about 1.10-1.40. The value y is determined by the proton NMR method.

R_1 in formula (I) is a C_8-18 linear or branched alkyl, alkenyl, or alkylphenyl group. From the aspect of 30 water-solubility and foaming ability, a preferable carbon number is C_{10-14} . R_2 in formula (I) is a C_2-4 alkylene group. From the aspect of water-solubility and the like, a preferable carbon number is C_{2-3} . The structure of G in formula (I) depends upon whether it represents a monosaccharide or a di- or higher saccharide. Monosaccharides include glucose, galactose, xylose, mannose, lyxose and arabinose, as well as their mixtures. Examples of di- or higher saccharides are maltose, xylobiose, isomaltose, cellobiose, 35 gentiobiose, lactose, sucrose, nigerose, turanose, raffinose, gentianose and melezitose, as well as their mixtures. Among these, from the aspect of availability and cost, preferable monosaccharides are glucose and fructose, and preferable di- or higher saccharides are maltose and sucrose.

Component (a) is incorporated in the composition of the present invention in an amount of 2-60%, preferably 10-40% by weight.

40 Component (b) of the present invention is nonionic surfactants having an HLB value of less than 5. The HLB value is defined as the value calculated from the equation, J. T. Davies and E. K. Rideal: *Interfacial Phenomena*, pp 371-383 (1963), Academic Press, New York.

$$HLB = 7 + \Sigma (\text{number of hydrophilic groups}) \\ - \Sigma (\text{number of hydrophobic groups})$$

The number of each atomic group used for the HLB calculation is as follows.

	Atomic group	number of groups
5	<hr/>	
	Hydrophilic group	
	Ester (sorbitan ring)	6.8
10	Ester (free)	2.4
	-COOH	2.1
15	OH (free)	1.3
	-O-	0.5
	OH (sorbitan ring)	0.5
20	<hr/>	
	Lipophilic group	
	$\begin{array}{c} \\ -CH- \end{array}$	0.475
25	-CH ₂ -	0.475
	CH ₃ -	0.475
30	=CH-	0.475
	<hr/>	
	Induction group	
35	-(CH ₂ -CH ₂ -O)-	0.33
	$\begin{array}{c} -(CH_2-CH-O)- \\ \\ CH_3 \end{array}$	0.15
40	<hr/>	

Examples of component (b) are polyoxyethylene alkyl ether, polyoxyethylene alkylphenyl ether, polyoxyethylene sorbitan fatty acid ester, polyoxyethylene sorbit fatty acid ester, polyoxyethylene glycol fatty acid ester, polyoxyethylene polyoxypropylene alkyl ether, polyoxyethylene castor oil, polyoxyethylene hydrogenated castor oil, polyoxyethylene alkylamine and glycerol fatty acid ester. Component (b) is incorporated in the composition of the present invention in an amount of 0.1-10%, preferably 0.5-8% by weight.

Component (c) of the present invention is nonionic surfactants having an HLB value of not less than 5. Enumerated as examples of component (c) are polyoxyethylene alkyl ether, polyoxyethylene alkylphenyl ether, polyoxyethylene sorbitan fatty acid ester, polyoxyethylene sorbit fatty acid ester, polyoxyethylene glycol fatty acid ester, polyoxyethylene polyoxypropylene alkyl ether, polyoxyethylene castor oil, polyoxyethylene hydrogenated castor oil, polyoxyethylene alkylamine and polyoxyethylene fatty acid amide, having the total adduct mols of ethylene oxide of not less than 10.

Component (c) is incorporated in the composition of the present invention in an amount of 0.1-10%, preferably 0.3-8% by weight.


The ratio of component [(b) + (c)]/(a) is in the range of 1/10-1/1, preferably 1/8-1/1.5 by weight. The ratio of component (b)/(c) is in the range of 1/10-1/1, preferably 1/8-1/1.5 by weight.

The difference in HLB between component (b) and component (c) is preferably greater than 3.

Component (d) used in the present invention is water-soluble organic or inorganic salt given in the aforementioned explanation. Among them, sulfates, chlorides, borates, p-toluenesulfonates, benzoates, citrates, and edates of sodium and potassium are particularly preferable.

The composition of the present invention contains 0.001-8% by weight of component (d). A preferable amount of component (d) is 0.01-5%, and particularly 0.1-4% by weight. The amount exceeding 8% by weight not only impairs the intended effects, but also adversely affects the stability of the composition.

Besides the above essential components, the composition of the present invention may comprise one or more of surfactants selected from the groups consisting of anionic surfactants (1-20% by weight), amphoteric surfactants (1-20% by weight), and nonionic surfactants (1-10% by weight) other than the above components (a)-(c). These surfactants are optionally incorporated into the composition of the present invention in order to promote its detergency and foaming ability. conventionally known anionic or amphoteric surfactants can be used. Such surfactants may include, for example, α -olefin sulfonates [C_8-20 ; Na, K, Mg, triethanolamine (TEA), NH_4 salts], polyoxyethylene alkylsulfates (EO = 2-8; alkyl group: C_8-18 , linear or branched; Na, K, Mg, TEA, NH_4 salts), salts of α -sulfo-fatty acid ester represented by the formula,



(wherein R_5 : C_8-18 , R_6 : C_1-4 , M: alkali metals, M-acylglutamates (acyl group: C_8-18 ; Na, K, TEA salts), monoalkyl phosphates (alkyl group: C_8-18 ; Na, K, TEA, arginine salts), linear alkylbenzenesulfonates (alkyl group: C_{10-18} ; Na, K, Mg salts), alkyl betaine, alkyl sulfobetaine and alkyl hydroxysulfobetaine. Given as examples of the nonionic surfactants other than component (a)-(c) are mono- or di-alkanol (C_2-3) amides of C_3-22 fatty acid, and tert-alkylamine oxides (alkyl group: C_8-18 , linear or branched).

The pH range of a neat liquid of the detergent composition of the present invention is preferably 6-8, particularly 6.5-7.5 is most preferable.

In addition to the above components, other optional ingredients can be formulated into the composition of the present invention to the extent that its stability, detergency, and foaming ability are not adversely affected. Such optional components include lower aliphatic alcohols, e.g. ethanol; solubilizers, e.g. urea; viscosity adjusting agents, e.g. clay minerals, water-soluble polymers; water-insoluble abrasives, e.g. calcite, silica, calcium phosphate, zeolite, polyethylene, nylon, polystyrene; moisturizing agents, e.g. glycerol, sorbitol; feeling improvers, e.g. cationized cellulose; enzymes; perfumes; pigments; antiseptics; and antifungal agents.

The detergent composition of the present invention thus prepared has excellent foaming ability, is mild to the skin and the hair, and imparts a favorable feeling to the hands during and after use. The foam is easily rinsed off and after washing water is easily drained from the tableware and the like. Thus, it is a neutral detergent composition having a high practical value.

Other features of the invention will become apparent in the course of the following description of the exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

EXAMPLES

Example 1

The detergent compositions having the formulations shown in Table 1 were prepared. Their foaming ability, rinsability, feeling to the hands during and after use, and drainability were evaluated. The results are shown in Table 1.

The feeling to the hands during and after use was tested using composition A (Composition 11 in Table 1) as a standard. Evaluation was made by comparing the feelings given to the hands when composition A and composition B (Compositions 1-10 in Table 1) were used.

(Test Methods and Evaluation Standards)

(1) Foaming ability

- 5 Foaming ability was measured on samples prepared by adding a commercially available butter in an amount of 0.1%, as a dirt component, to 0.5% detergent composition solutions. Forty milliliters of the solution of the detergent composition to which butter was added was charged into a glass cylinder with a diameter of 5 cm and subjected to a rotational stirring for 10 minutes. The height of the foam produced was measured after the termination of the stirring.

10

(2) Rinsability

- 15 Three liters of a sample solution with 0.25% detergent concentration was charged into a container (diameter: 30 cm, height: 12 cm) and rotationally stirred for 10 minutes, after which the liquid was discharged through an opening at the bottom of the container, leaving the foam in the container. Three liters of tap water was then charged to the container, and the ten-minutes stirring and discharge of liquid were repeated. This procedure was repeated until no foam was observed in the container. The number of times of tap water charge and discharge required for the foam to diminish was taken as the standard of rinsability.

20 (3) Feeling to the hands during and after use

Two 10% detergent composition solutions, A and B, at 40 °C were charged into 2-liter beakers. Feeling of solutions A and B to the hands was sensuously evaluated according to the following standard.

25 (i) Feeling of the solution

Hands were dipped into solutions A and B, i.e., one hand to solution A and the other to solution B. After 1 minute, the feeling to the hands was compared according to the following standard.

30

B is less slimy than A	+2
B is slightly less slimy than A	+1
Hard to tell which is more slimy	0
B is slightly more slimy than A	-1
B is more slimy than A	-2

35

(ii) Feeling after use

- 40 After the above test, the hands were thoroughly washed with water and wiped out with a dry towel. Then feeling of the hands were compared according to the following standard.

45

B is less sticky than A	+2
B is slightly less sticky than A	+1
Hard to tell which is more sticky	0
B is slightly more sticky than A	-1
B is more sticky than A	-2

- 50 The above tests were carried out by 10 panelists. The total scores gained by solution B was taken as the feeling of the solution B.

(4) Drainability

- 55 Three liters of a 0.15 % detergent composition solution (in 3.5° DH at 25 °C) was charged into a container (diameter: 30 cm, depth: 12 cm). A piece of porcelain dish having a 25 cm diameter was dipped into the solution for 3 minutes. The dish was taken out from the solution and dried at 25 °C. This procedure was repeated 4 times. Then the dish was again dipped into the solution, following which it was taken out

from the solution and rinsed for 30 seconds in a water stream and dried. The number of water spots remaining on the surface of the dish was counted to evaluate the drainability according to the following standard.

5 Evaluation Standard:

- A: No water spots are present
 B: Less than 20 water spots are present
 C: 20 or more of water spots are present

10

TABLE 1

Component	Example Compositions										z by weight
	1	2	3	4	5	6	7	8	9	10	11
<u>Component (a)</u>											
Alkyl glycoside: in formula (I), x=0, C-glucose residue,											
R ₁ : C ₉₋₁₁ alkyl, γ=1.2	23	23	-	-	23	-	23	23	-	-	23
R ₁ : C ₉₋₁₁ alkyl, γ=1.42	-	-	23	23	-	23	-	-	23	23	-
<u>Component (b)</u>											
Polyoxyethylene(p=3) lauryl ether (HLB-2.8)											
	3	-	-	4	-	10	3	-	-	-	-
Glycerol monooleate (HLB-2.5)											
		3	-	-	1	-	-	-	-	1	0.05
Sorbitan monostearate (HLB-4.9)											
	-	-	1	-	-	-	-	-	3	-	-
<u>Component (c)</u>											
Polyoxyethylene(p=15) lauryl ether (HLB-6.8)											
	5	-	-	4	-	-	-	-	-	-	-
Polyethylene glycol (HM-3000) monostearate (HLB-23.7)											
	-	5	-	-	10	-	-	5	-	-	0.05
Polyoxyethylene(p=20) polyoxypropylene(p=5) syringyl ether (HLB-8.2)											
	-	-	2	-	-	-	-	-	-	12	-
Polyoxyethylene(p=20) sorbitan monolaurate (HLB-14.3)											
	-	-	-	-	-	10	-	-	5	-	-

TABLE 1 (Continued)

Component	Example Compositions						Comparative Compositions				
	1	2	3	4	5	6	7	8	9	10	11
<u>Component (d)</u>											
NaCl	0.5	-	-	-	-	-	0.5	-	-	-	-
K ₂ SO ₄	-	0.5	-	0.2	-	-	-	-	-	-	0.5
Sodium citrate	-	-	0.5	-	-	3	-	0.5	-	-	-
Sodium p-toluenesulfonate	-	-	-	0.3	-	-	-	-	-	-	-
Disodium edate	-	-	-	-	0.5	-	-	-	-	0.5	-
Water	Balance	Balance	Balance	Balance	Balance	Balance	Balance	Balance	Balance	Balance	Balance
pH 1) (neat liquid)	6.8	7.0	7.1	6.3	7.8	7.0	6.9	7.2	7.5	7.0	6.7
<u>Evaluation</u>											
Foaming ability (mm)	92	90	87	90	88	85	80	82	87	75	79
Rinsability (times)	5	5	5	5	5	5	7	7	7	8	7
Feeling to the hands											
During use	+9	+8	+8	+9	+8	+8	+2.0	+1.6	+3.5	+1.5	+0
After use	+9	+8	+8	+9	+8	+8	+2.3	+1.5	+3.5	+1.5	+0
Drainability	A	A	A	A	A	A	C	C	C	C	C

1) pH was appropriately adjusted with H₂SO₄ or NaOH.

2) p indicates the average moles of addition.

Example 2

5

10

15

Light duty liquid detergent composition for laundry	
	% by weight
Alkyl glycoside 1)	20
Sodium polyoxyethylene($\bar{p}=2$)alkyl(C_{12-13})ether sulfate	10
Softhanol 33 2)	2
Polyoxyethylene($\bar{p}=30$)lauryl ether (HLB = 11.7)	3
Sodium malate	3
Ethanol	4
Perfume, enzyme	Small amount
Water	Balance
	100
(The above mixture was adjusted to pH 8.0)	

20

1) In formula (I), $x=1$, $y=1.30$, $R_1=C_{10-12}$ alkyl, $R_2=C_2$ alkylene, G = glucose residue

2) Trademark, a nonionic surfactant produced by Nippon Shokubai Kagaku Kogyo Co., Ltd. (HLB: about 2.4)

25 Example 3

30

35

40


Dishwashing detergent composition	
	% by weight
Alkyl glycoside 1)	20
Sodium polyoxyethylene($\bar{p}=5$)laurylether sulfate	5
Glycerol monomyristate (HLB = 4.4)	2
Polyoxyethylene($\bar{p}=60$)hydrogenated castor oil	5
Lauryldimethylamine oxide	2
Sodium sulfate	1
Ethanol	3
Perfume, enzyme	Small amount
Water	Balance
	100
(The above mixture was adjusted to pH 7.0)	

45

1) In formula (I), $x=0$, $y=1.4$, $R_1=C_9-11$ alkyl, G = glucose residue

Claims

1. A neutral liquid detergent composition comprising:
 (a) 2-60% by weight of an alkyl glycoside represented by the following formula (I):



wherein R_1 is a linear or branched alkyl, alkenyl, or alkylphenyl group having 8-18 carbon atoms, R_2 is an alkylene group having 2-4 carbon atoms, G is a reducing saccharide residue having 5-6 carbon atoms, x is a number of which the mean value is 0-5, and y is a number of which the mean value is 1.0-1.42;

- (b) 0.1-10% by weight of a nonionic surfactant having an HLB of less than 5;

- (c) 0.1-10% by weight of a nonionic surfactant having an HLB of not less than 5; and
 (d) 0.001-8% by weight of one or more water-soluble organic or inorganic salts selected from the group consisting of sulfates, chlorides, borates, phosphates, p-toluensulfonates, m-xylenesulfonates, benzoates, malates, succinates, tartarates, citrates, lactates, and edates of sodium and potassium and mixtures thereof; the ratio by weight of component [(b) + (c)]/(a) being in a range of 1/10-1/1 and the ratio by weight of component (b)/(c) being in a range of 1/10-1/1.
- 5
2. A neutral liquid detergent composition according to Claim 1, further comprising one or more surfactants selected from 1-20% by weight of anionic surfactants, 1-20% by weight of amphoteric surfactants, and 1-10% by weight of nonionic surfactants other than components (a)-(c).
- 10

Patentansprüche

1. Flüssiges Neutralwaschmittel, umfassend:
- 15 (a) 2 bis 60 Gew.-% eines Alkylglycosids, wiedergegeben durch die nachstehende Formel (I):
- $$R_1(OR_2)_xG_y \quad (I),$$
- worin R_1 eine lineare oder verzweigte Alkyl-, Alkenyl- oder Alkylphenylgruppe mit 8 bis 18 Kohlenstoffatomen bedeutet, R_2 eine Alkylengruppe mit 2 bis 4 Kohlenstoffatomen bedeutet, G einen reduzierenden Saccharidrest mit 5 bis 6 Kohlenstoffatomen bedeutet, x eine Zahl ist, deren Mittelwert 0 bis 5 ist, und y eine Zahl ist, deren Mittelwert 1,0 bis 1,42 ist;
- 20 (b) 0,1 bis 10 Gew.-% eines nichtionischen Tensids mit einem HLB-Wert von weniger als 5;
- (c) 0,1 bis 10 Gew.-% eines nichtionischen Tensids mit einem HLB-Wert von nicht weniger als 5;
- 25 und
- (d) 0,001 bis 8 Gew.-% von einem oder mehreren wasserlöslichen organischen oder anorganischen Salzen, ausgewählt aus der Gruppe, bestehend aus Sulfaten, Chloriden, Boraten, Phosphaten, p-Toluolsulfonaten, m-Xylolsulfonaten, Benzoaten, Malaten, Succinaten, Tartraten, Citraten, Lactaten und Edaten von Natrium und Kalium und Gemischen davon; wobei das Gewichtsverhältnis der Komponente [(b) + (c)]/(a) in einem Bereich von 1/10 bis 1/1 ist und das Gewichtsverhältnis der Komponente (b)/(c) in einem Bereich von 1/10 bis 1/1 ist.
- 30
2. Flüssiges Neutralwaschmittel nach Anspruch 1, weiterhin umfassend eines oder mehrere Tenside, ausgewählt aus 1 bis 20 Gew.-% anionischer Tenside, 1 bis 20 Gew.-% amphoterer Tenside und 1 bis 10 Gew.-% nichtionischer Tenside, die von Bestandteilen (a) - (c) verschieden sind.
- 35

Revendications

1. Composition détergente liquide neutre, comprenant :
- 40 (a) 2 - 60% en poids d'un alkyl glycoside, représenté par la formule suivante (I) :
- $$R_1(OR_2)_xG_y \quad (I)$$
- dans laquelle :
- 45 - R_1 représente un groupe alkyle, alcényle ou alkylphényle, linéaire ou ramifié, ayant 8 - 18 atomes de carbone ;
- R_2 représente un groupe alkylène ayant 2 - 4 atomes de carbone ;
- G représente un résidu de saccharide réducteur ayant 5 - 6 atomes de carbone ;
- x est un nombre dont la valeur moyenne est 0 - 5 ; et
- 50 - y est un nombre dont la valeur moyenne est 1,0 - 1,42 ;
- (b) 0,1 - 10% en poids d'un agent tensio-actif non-ionique ayant une valeur HLB de moins de 5 ;
- (c) 0,1 - 10% en poids d'un agent tensio-actif non-ionique ayant une valeur HLB non inférieure à 5 ; et
- (d) 0,001 - 8% en poids d'un ou plusieurs sels organiques ou minéraux, solubles dans l'eau, choisis dans le groupe constitué par les sulfates, chlorures, borates, phosphates, p-toluène sulfonates, m-xylène sulfonates, benzoates, malates, succinates, tartrates, citrates, lactates et édétates de sodium et de potassium, et leurs mélanges ;
- 55 le rapport en poids des composants [(b) + (c)]/(a) se situant dans une plage de 1/10 - 1/1 et le

rapport en poids des composants (b)/(c) se situant dans une plage de 1/10 - 1/1.

2. Composition détergente liquide neutre selon la revendication 1, comprenant en outre un ou plusieurs agents tensio-actifs choisis parmi 1 - 20% en poids d'agents tensio-actifs anioniques, 1 - 20% d'agents tensio-actifs amphotères, et 1 - 10% en poids d'agents tensio-actifs non-ioniques autres que les composants (a)-(c).

10

15

20

25

30

35

40

45

50

55